

Constraining CP violation in a softly broken A_4 symmetric Model

Biswajit Adhikary* and Ambar Ghosal†

Saha Institute of Nuclear Physics,
1/AF Bidhan Nagar, Kolkata 700064, India

Abstract

To understand the mass spectra of charged lepton and neutrino A_4 symmetry has been proposed in addition with the Standard $SU(2)_L \times U(1)_Y$ model. We break A_4 symmetry softly and the deviation from the tri-bimaximal mixing arises due to Zee mechanism. In the present work, we express two mixing angles θ_{13} and θ_{23} in terms of a single model parameter and experimental observables, such as, mixing angle θ_{12} , mass squared differences Δm_{21}^2 and Δm_{32}^2 . Using the experimental values of θ_{23} , θ_{12} , Δm_{21}^2 and Δm_{32}^2 we restrict the model parameter and we predict θ_{13} . This model gives rise to $\theta_{13} \simeq 11^\circ$ if we allow 1σ deviation of θ_{23} and 2° deviation of θ_{12} from their best fit values. Utilizing all those constraints, we explore the extent of CP violation parameter J_{CP} in the present model and found a value of $J_{CP} \approx 2.65 \times 10^{-3}$ (for 1σ deviation of θ_{23} and 2° deviation of θ_{12}) consistent with the other neutrino experimental results. We have studied the mass pattern of neutrino and neutrinoless double beta decay ($\beta\beta_{0\nu}$) parameter $|(M_\nu)_{ee}|$ in this model.

*biswajit.adhikary@saha.ac.in

†ambar.ghosal@saha.ac.in

An interesting way to obtain hierarchical charged lepton mass matrix along with appropriate texture of neutrino mass matrix which can accommodate present neutrino experimental results, namely, solar, atmospheric and CHOOZ is through the introduction of non-abelian discrete A_4 symmetry in a model [1]. The interplay of A_4 symmetry predicts diagonal and hierarchical charged lepton mass matrix in addition with the neutrino masses which could be quasi-degenerate or hierarchical. Altarelli and Feruglio have proposed a version of A_4 symmetric model (AF model) [2] to obtain hierarchical charged lepton mass matrix along with tri-bimaximal neutrino mixing ($\sin \theta_{12} = 1/\sqrt{3}$, $\sin \theta_{23} = -1/\sqrt{2}$, $\sin \theta_{13} = 0$) [3]. Although the model gives rise to $\theta_{13} = 0$ ($|U_{e3}| = 0$) which is consistent with the CHOOZ-Palo Verde experimental upper bound ($\theta_{13} < 12^\circ$ at 3σ), however, the non-zero value of $|U_{e3}|$ opens up a possibility to explore CP violation in leptonic sector which is the main goal of many future short and long baseline experiments.

A prediction for non-zero U_{e3} has been realized in a recently proposed modified AF model through the inclusion of three gauge singlet charged scalars due to radiative correction of the off diagonal elements of the neutrino mass matrix [4]. The model successfully predicts solar and atmospheric neutrino mixing and mass-squared differences along with small but non-zero value of θ_{13} well below the present experimental upper bound for a reasonable choice of model parameters. A relationship between different mixing angles is an outcome of the model and the predictability of the model is also testable in future neutrino experiments.

Aim of this paper is to generalize the assumptions made in [4]. In the present work we bring down the value of θ_{12} to its best fit value $\theta_{12} = 34.0^\circ$ from the tri-bimaximal value of the above $\theta_{12} = 35.26^\circ$ which is at the 1σ edge of experimental value. We investigate how much θ_{13} can be within 1σ variation of θ_{23} and at the best fit value as well as 2° variation about the best fit value of θ_{12} . Then we shift our concentration to CP violating parameter J_{CP} and Dirac phase δ_D and figure out their values. We also see the mass pattern of the neutrinos and also see the variation of $\beta\beta_{0\nu}$ experimental parameter in this model. For our analysis we have used the best fit value of the mass squared differences, $\Delta m_{21}^2 = 8.0 \times 10^{-5} \text{ eV}^2$ and $\Delta m_{32}^2 = 2.1 \times 10^{-3} \text{ eV}^2$.

For completeness, we briefly summarize here the model proposed in [4]. The lepton content with their representation and the Higgs content with their vevs and representation under $SU(2)_L \times U(1)_Y \times A_4$ symmetry is presented in Table 1, where all fields except the charged scalar χ_i^+ have been used in the original AF model [2]. The Yukawa interaction in the leptonic sector is given by

$$L_{AF}^l = y_e e^c (\phi_T l) h_d / \Lambda + y_\mu \mu^c (\phi_T l)' h_d / \Lambda + y_\tau \tau^c (\phi_T l)'' h_d / \Lambda +$$

$$x_a \xi(lh_u lh_u)/\Lambda^2 + x_b(\phi_S lh_u lh_u)/\Lambda^2 \quad (1)$$

where $x_a, x_b, y_e, y_\mu, y_\tau$ are Yukawa couplings and Λ is the new mass scale. After spontaneous breaking of the symmetry of the model, the above Lagrangian gives rise to the following mass terms

$$\begin{aligned} \mathcal{L}_{AF} = & v_d v_T / \Lambda (y_e e^c e + y_\mu \mu^c \mu + y_\tau \tau^c \tau) \\ & + x_a v_u^2 (u/\Lambda^2) (\nu_e \nu_e + 2\nu_\mu \nu_\tau) \\ & + x_b v_u^2 2v_S / 3\Lambda^2 (\nu_e \nu_e + \nu_\mu \nu_\mu \\ & + \nu_\tau \nu_\tau - \nu_e \nu_\mu - \nu_\mu \nu_\tau - \nu_\tau \nu_e) + h.c. \end{aligned} \quad (2)$$

The charged lepton and neutrino mass matrices come out as

$$m_l = \begin{pmatrix} m_e & 0 & 0 \\ 0 & m_\mu & 0 \\ 0 & 0 & m_\tau \end{pmatrix}, m_\nu^{\text{AF}} = m_0 \begin{pmatrix} a + 2d/3 & -d/3 & -d/3 \\ -d/3 & 2d/3 & a - d/3 \\ -d/3 & a - d/3 & 2d/3 \end{pmatrix}, \quad (3)$$

where

$$\begin{aligned} m_e = & y_e v_d v_T / \Lambda, m_\mu = y_\mu v_d v_T / \Lambda, m_\tau = y_\tau v_d v_T / \Lambda, \\ a = & 2x_a u / \Lambda, d = 2x_b v_S / \Lambda, m_0 = v_u^2 / \Lambda. \end{aligned} \quad (4)$$

The charged lepton mass matrix is diagonal so the leptonic mixing solely occurs from the neutrino sector and diagonalising m_ν by the way $U^\dagger m_\nu U^*$ we get the three mass eigenvalues as

$$m_1 = a + d, m_2 = a, m_3 = d - a \quad (5)$$

with the exact tri-bimaximal mixing pattern. Thus, in the AF model, tri-bimaximal mixing occurs naturally. In [4], a modified version of the above model has been investigated to generate non-zero θ_{13} which at the leading order gives tri-bimaximal mixing form. In order to do that, an A_4 triplet $SU(2)_L$ singlet charged scalar $\chi_i^+ (= \chi_1^+, \chi_2^+, \chi_3^+)$ has been incorporated and the leptonic part of the Lagrangian becomes

$$L = L_{AF}^l + L_{MAF}^l \quad (6)$$

where L_{MAF}^l has two parts as

$$L_{MAF}^l = \mathcal{L}_1 + \mathcal{L}_2 \quad (7)$$

$\mathcal{L}_1$ is A_4 symmetry preserving part and is given by

$$\begin{aligned}\mathcal{L}_1 &= f (L L \chi_i) \subset (3 \times 3 \times 3) \\ &= f(\nu_\mu \tau \chi_1^+ + \nu_\tau e \chi_2^+ + \nu_e \mu \chi_3^+ - \nu_\tau \mu \chi_1^+ - \nu_e \tau \chi_2^+ - \nu_\mu e \chi_3^+).\end{aligned}\quad (8)$$

however, $\mathcal{L}_2$ contains Zee-type term which is explicit soft A_4 symmetry breaking and is given by

$$\mathcal{L}_2 = c_{12} h_u^T i \tau_2 h_d (\chi_1^+ + \chi_2^+ + \chi_3^+). \quad (9)$$

The charged lepton mass matrix is still diagonal in the present model and the neutrino mass matrix comes out as

$$m_\nu = \begin{pmatrix} a + 2d/3 & -d/3 & -d/3 - \epsilon \\ -d/3 & 2d/3 & a - d/3 + \epsilon \\ -d/3 - \epsilon & a - d/3 + \epsilon & 2d/3 \end{pmatrix} \quad (10)$$

where the a and d parameters defined earlier and are obtained due to higher dimensional operators in the same way as AF model. The ϵ term is the additional contribution arises at the one-loop level due to well known Zee mechanism and is shown in Fig.1. The parameter ϵ is given by

$$\epsilon = f m_\tau^2 \frac{c_{12} v_u}{v_d} F(m_\chi^2, m_{h_d}^2) \quad (11)$$

with the definition,

$$F(M_1^2, M_2^2) = \frac{1}{16\pi^2(M_1^2 - M_2^2)} \ln \frac{M_1^2}{M_2^2}. \quad (12)$$

Although the Lagrangian given in Eq.8 can generate corrections to all off-diagonal entries of the mass matrix given in Eq. (10), however, dominant terms proportional to m_τ^2 are retained. In this model neutrinos are Majorana-type in nature. In general, the parameters a , d , ϵ are all complex, however, it is possible to rotate out one of the phase. For our analysis, we consider only the parameter d is complex and parameters a, ϵ are real. The neutrino mass matrix in this case comes out as

$$M_\nu = \begin{pmatrix} a + 2de^{i\phi}/3 & -de^{i\phi}/3 & -de^{i\phi}/3 - \epsilon \\ -de^{i\phi}/3 & 2de^{i\phi}/3 & a - de^{i\phi}/3 + \epsilon \\ -de^{i\phi}/3 - \epsilon & a - de^{i\phi}/3 + \epsilon & 2de^{i\phi}/3 \end{pmatrix}. \quad (13)$$

A relationship between the parameters a and d has been considered as

$$d = \kappa a \cos \phi \quad (14)$$




Figure 1: One loop radiative $\nu_{e,\mu}$ - ν_τ mass due to charged Higgs exchange.

Lepton	$SU(2)_L$	A_4	
(ν_i, l_i)	2	3	
l_i^c	1	1	
Scalar			VEV
h_u	2	1	$\langle h_u^0 \rangle = v_u$
h_d	2	1	$\langle h_d^0 \rangle = v_d$
ξ	1	1	$\langle \xi^0 \rangle = u$
ϕ_S	1	3	$\langle \phi_S^0 \rangle = (v_S, v_S, v_S)$
ϕ_T	1	3	$\langle \phi_T \rangle = (v_T, 0, 0)$
χ_i^+	1	3	

Table 1: List of fermion and scalar fields used in this model.

where κ , d and a are the real parameters. The analysis of Ref.[4] has been done for a specific value of κ and in the present work, we have generalized the whole analysis. We constrain the value of κ first from the existing bounds of two mixing angles θ_{12} and θ_{23} . Then we have utilized the result to calculate θ_{13} and the CP violation parameter J_{CP} [5] and explore the extent at which J_{CP} is allowed in the present model.

With only assumption ϵ is small, we diagonalize mass matrix Eq. (13) as

$$U^\dagger M_\nu U^* = \text{diag} \left(de^{i\phi} + a + \epsilon, \quad a, \quad de^{i\phi} - a - \epsilon \right) \quad (15)$$

upto first order in ϵ where diagonalizing matrix

$$U = \begin{pmatrix} \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & -\sqrt{\frac{1}{2}} \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \end{pmatrix} + \epsilon \begin{pmatrix} -\frac{2a+de^{-i\phi}}{\sqrt{6}d(d+2a\cos\phi)} & \frac{2a+de^{i\phi}}{\sqrt{3}d(d+2a\cos\phi)} & \frac{1}{3\sqrt{2}} \left(\frac{1}{a} - \frac{e^{-i\phi}}{d-2a\cos\phi} \right) \\ \frac{d^2-4a^2+2iad\sin\phi}{2\sqrt{6}ad(d+2a\cos\phi)} & \frac{2a^2\cos\phi-d(a+de^{i\phi})}{\sqrt{3}(d^3-4a^2d\cos^2\phi)} & -\frac{1}{6\sqrt{2}} \left(\frac{1}{a} + \frac{2e^{-i\phi}}{d-2a\cos\phi} \right) \\ -\frac{4a^2+d^2+4ad\cos\phi-2iad\sin\phi}{2\sqrt{6}ad(d+2a\cos\phi)} & \frac{a(2a\cos\phi+de^{2i\phi})}{\sqrt{3}(d^3-4a^2d\cos^2\phi)} & -\frac{1}{6\sqrt{2}} \left(\frac{1}{a} + \frac{2e^{-i\phi}}{d-2a\cos\phi} \right) \end{pmatrix}. \quad (16)$$

It is to be noted that for vanishing value of ϵ the matrix U leads to tri-bimaximal form. Three approximate mass eigenvalues in Eq. (15) take the following forms

$$\begin{aligned} m_1^2 &= \left| de^{i\phi} + a + \epsilon \right|^2 \\ &\simeq a^2 \left[1 + 2\epsilon' + 2\kappa \cos^2 \phi + 2\epsilon' \kappa \cos^2 \phi + \kappa^2 \cos^2 \phi \right] \\ m_2^2 &= a^2 \\ m_3^2 &= \left| de^{i\phi} - a - \epsilon \right|^2 \\ &\simeq a^2 \left[1 + 2\epsilon' - 2\kappa \cos^2 \phi - 2\epsilon' \kappa \cos^2 \phi + \kappa^2 \cos^2 \phi \right] \end{aligned} \quad (17)$$

where $\epsilon' = \epsilon/a$ and we use the relation given in Eq. (14). The solar and atmospheric neutrino mass squared differences are coming out as

$$\begin{aligned} \Delta m_{\odot}^2 &= \Delta m_{21}^2 = m_2^2 - m_1^2 = a^2 \left[-\kappa \cos^2 \phi (\kappa + 2) - 2\epsilon' (1 + \kappa \cos^2 \phi) \right] \\ \Delta m_{\text{atm}}^2 &= \Delta m_{32}^2 = m_3^2 - m_2^2 = a^2 \left[-\kappa \cos^2 \phi (2 - \kappa) + 2\epsilon' (1 - \kappa \cos^2 \phi) \right]. \end{aligned} \quad (18)$$

From mixing matrix U in Eq. (16) we obtain the mixing angles:

$$\sin \theta_{12} = |U_{12}| = \frac{1}{\sqrt{3}} + \frac{\epsilon' (2 + \kappa \cos^2 \phi)}{\sqrt{3} \kappa \cos^2 \phi (\kappa + 2)}$$




Figure 2: Plot of $\cos^2 \phi$ with respect to κ and θ_{12} . We keep Δm_{32}^2 and Δm_{21}^2 to their best fit values.

$$\begin{aligned} \sin \theta_{13} = |U_{13}| &= \left| \frac{\epsilon'}{3\sqrt{2}(\kappa - 2)} \right| \left[\frac{1 + \cos^2 \phi (\kappa^2 + 8 - 6\kappa)}{\cos^2 \phi} \right]^{1/2} \\ \tan^2 \theta_{23} &= \frac{|U_{23}|^2}{1 - |U_{23}|^2} = 1 + \frac{2\epsilon'\kappa}{3(\kappa - 2)}. \end{aligned} \quad (19)$$

We define the ratio R as

$$R = \frac{\Delta m_{\odot}^2}{\Delta m_{atm}^2} = \frac{\kappa \cos^2 \phi (\kappa + 2) + 2\epsilon'(1 + \kappa \cos^2 \phi)}{\kappa \cos^2 \phi (2 - \kappa) - 2\epsilon'(1 - \kappa \cos^2 \phi)}. \quad (20)$$

Using the first relation of Eq. (19) we have

$$\epsilon' = \frac{\sqrt{3}\kappa \cos^2 \phi (\kappa + 2)}{2 + \kappa \cos^2 \phi} \times (\sin \theta_{12} - 1/\sqrt{3}). \quad (21)$$

Using Eq. (21) in Eq. (20) we get

$$\cos^2 \phi = \frac{-2}{\kappa} \left[\frac{2(1 - R) + \kappa(1 + R) + \sqrt{3}(\kappa + 2)(\sin \theta_{12} - 1/\sqrt{3})(1 + R)}{2(1 - R) + \kappa(1 + R) + 2\sqrt{3}(\kappa + 2)(\sin \theta_{12} - 1/\sqrt{3})(1 - R)} \right]. \quad (22)$$

Again using expression of ϵ' from Eq. (21) to the expression of Δm_{32}^2 in Eq. (18) we get the dependence of the a^2 on κ and $\cos^2 \phi$:

$$a^2 = \frac{\Delta m_{32}^2 (2 + \kappa \cos^2 \phi)}{\kappa \cos^2 \phi [(\kappa - 2)(2 + \kappa \cos^2 \phi) + 2\sqrt{3}(\kappa + 2)(1 - \kappa \cos^2 \phi)(\sin \theta_{12} - 1/\sqrt{3})]} \quad (23)$$




Figure 3: Plot of θ_{23} with respect to κ and θ_{12} . We keep Δm_{32}^2 and Δm_{21}^2 to their best fit values.

Thus from Eq. (22) and Eq. (23), we see that a^2 only depends on single model parameter κ . From the above expression of $\cos^2 \phi$ we can put bound on κ for the given values of R and θ_{12} . For the best fit value of Δm_{32}^2 and Δm_{21}^2 we have the approximate bound:

$$\kappa < -2 \quad (24)$$

in the course of variation $32^\circ \leq \theta_{12} \leq 36^\circ$ for $\cos^2 \phi < 1$ and it is shown in Fig. 2. It also ensures $\cos^2 \phi > 0$. Again compatibility of the above bound of κ with the restriction $a^2 > 0$ demands that $\Delta m_{32}^2 > 0$. This leads to normal ordering of neutrino masses.

Using the expression of ϵ' from Eq. (21) into the expression of $\sin \theta_{13}$ and $\tan^2 \theta_{23}$ in Eq. (19) we get the two mixing angles in terms of κ , $\cos^2 \phi$:

$$\tan^2 \theta_{23} = 1 + \frac{2\sqrt{3}\kappa^2 \cos^2 \phi (\kappa + 2)}{3(\kappa - 2)(2 + \kappa \cos^2 \phi)} \times (\sin \theta_{12} - 1/\sqrt{3}) \quad (25)$$

and

$$\sin \theta_{13} = \left| \frac{\kappa \cos^2 \phi (\kappa + 2)}{\sqrt{6}(\kappa - 2)(2 + \kappa \cos^2 \phi)} \right| \left[\frac{1 + \cos^2 \phi (\kappa^2 + 8 - 6\kappa)}{\cos^2 \phi} \right]^{1/2} \left| \sin \theta_{12} - 1/\sqrt{3} \right| \quad (26)$$

Here also $\tan^2 \theta_{23}$ and $\sin \theta_{13}$ only depend on κ as $\cos^2 \phi$ is only function of κ . In addition to this dependence on parameter κ , they also depend on the well measured quantities, mixing angle θ_{12} , and the ratio of solar and atmospheric mass-squared differences R . We keep the solar and atmospheric mass-squared differences on their best fit values. We study the variation of θ_{23} with mixing angle θ_{12} and the parameter κ in Fig.3. We have varied κ from its analytical upper bound -2.0 to -2.5 and also have varied θ_{12} from 32° to 36° (2° deviation about best fit value 34° of θ_{12}). For a fixed value of κ variation of θ_{23} with θ_{12} is small and is smaller in higher value of κ , e.g. for the variation of θ_{12} from 32° to 36° for $\kappa = -2.0$, θ_{23} remains almost at 45.8° and for $\kappa = -2.5$, θ_{23} changes from 48.4° to 48.6° . If we allow θ_{23} upto its 1° deviated value 46° , range of κ shrinks to $-2.04 \leq \kappa \leq -2.0$ which is insensitive to the variation of θ_{12} . To keep θ_{23} within 1σ deviated value 48° for the whole range of variation of θ_{12} from 32° to 36° , we have to keep $\kappa \geq -2.39$. From Fig.3 we can also study the variation of θ_{23} with κ for a fixed value of θ_{12} , e.g. for the best fit value of $\theta_{12} = 34^\circ$, θ_{23} changes from 45.8° to 48.5° for the variation of κ from -2.0 to -2.5 . So, from Fig.3 we have gathered the information where we should keep the value of κ in the light of experimental values of θ_{23} and θ_{12} . With this information we study other observables.

Fig.4, we have plotted θ_{13} with θ_{12} and κ in their allowed region using the best fit value of solar and atmospheric mass squared differences. We have promised to generate nonzero θ_{13} with changing θ_{12} from its tri-bimaximal value 35.26° . Thus, we obtain the prediction on θ_{13} as $2.91^\circ \leq \theta_{13} \leq 3.7^\circ$ for $-2.04 \leq \kappa \leq -2.0$ (alternatively for $45.8^\circ \leq \theta_{23} \leq 46.0^\circ$) for the whole range of variation of θ_{12} 32° to 36° . If we allow θ_{23} upto its 1σ deviated value $45.8^\circ \leq \theta_{23} \leq 48.0^\circ$ (equivalently $-2.39 \leq \kappa \leq -2.0$), we have the prediction of θ_{13} as $2.91^\circ \leq \theta_{13} \leq 10.7^\circ$ for the whole range of variation of θ_{12} from 32° to 36° . So, the upper bound is near the largest possible allowed value from the experiment (CHOOZ $\theta_{13} < 12^\circ$ at 3σ). From the plot of Fig.4, this is to be noted that the upper bound of θ_{13} is mildly sensitive to θ_{12} , it varies from 9.93° to 10.7° at $\kappa = -2.39$. But the lower bound of θ_{13} is 2.91° which is insensitive to the variation of θ_{12} at $\kappa = -2.0$.

Keeping all those constraints in view next we explore the parameter space of CP violation parameter J_{CP} . The parameter J_{CP} defined as [5]

$$J_{CP} = \frac{1}{8} \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \sin \delta_D = \frac{\text{Im}[h_{12}h_{23}h_{31}]}{\Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2} \quad (27)$$




Figure 4: Plot of θ_{13} with respect to κ and θ_{12} . We keep Δm_{32}^2 and Δm_{21}^2 to their best fit values.

where $h = M_\nu M_\nu^\dagger$, δ_D is Dirac phase. This J_{CP} is associated with CP violation in neutrino oscillation and is directly related to Dirac phase of mixing matrix. Using Eq. (13), Eq. (14) in Eq. (27) CP violating parameter J_{CP} takes the following form

$$J_{CP} = \frac{a^6}{\Delta m_{21}^2 \Delta m_{32}^2 \Delta m_{31}^2} \times \frac{2\epsilon'}{9} \kappa^3 (\kappa + 2) \cos^4 \phi \sqrt{\cos^2 \phi - \cos^4 \phi}. \quad (28)$$

upto first order term in ϵ' . Using the expression of ϵ' from Eq. (21) and a^2 from Eq. (23) into the Eq. (28) we have

$$J_{CP} = \frac{2\sqrt{3}}{9} \frac{\kappa(\kappa + 2)^2 (2 + \kappa \cos^2 \phi)^2 (\sin \theta_{12} - 1/\sqrt{3}) \sqrt{\cos^2 \phi - \cos^4 \phi}}{R[(\kappa - 2)(2 + \kappa \cos^2 \phi) + 2\sqrt{3}(\kappa + 2)(1 - \kappa \cos^2 \phi)(\sin \theta_{12} - 1/\sqrt{3})]^3}. \quad (29)$$

J_{CP} is also only function of κ because $\cos^2 \phi$ in Eq. (22) is function of κ only. It is clear from the above expression if $\kappa = -2$ the value of J_{CP} becomes zero. This $\kappa + 2$ factor appear in J_{CP} in Eq. (28) is purely from A_4 symmetric structure of the neutrino mass matrix. In our analysis we have seen that κ and $\cos^2 \phi$ are not independent. $\kappa = -2$ approximately corresponds to $\cos^2 \phi = 1$ where J_{CP} also vanishes. Using the expression of $\cos^2 \phi$ from Eq. (22) into above Eq. (29) we can make J_{CP} only κ dependent. It also depends on the observables quantities θ_{12} , Δm_{32}^2 and Δm_{21}^2 . We have plotted in Fig.5 J_{CP} with respect κ and θ_{12} in their allowed region keeping Δm_{32}^2 and Δm_{21}^2 to their best fit values. We have the values of J_{CP} in the range $3.27 \times 10^{-5} \leq J_{CP} \leq 7.6 \times 10^{-5}$ for $-2.04 \leq \kappa \leq -2.0$ (alternatively for $45.8^\circ \leq \theta_{23} \leq 46.0^\circ$) for the whole range of variation of θ_{12} from 32° to 36° . If we allow θ_{23} upto its 1σ deviated value $45.8^\circ \leq \theta_{23} \leq 48.0^\circ$ (equivalently $-2.39 \leq \kappa \leq -2.0$), we have the prediction of J_{CP} as $3.27 \times 10^{-5} \leq J_{CP} \leq 2.65 \times 10^{-3}$ for the whole range of variation of θ_{12} from 32° to 36° . Upper bound of J_{CP} is varying from 2.12×10^{-3} to 2.65×10^{-3} for the variation of θ_{12} from 32° to 36° . For the best fit value of θ_{12} ($\approx 34^\circ$), upper bound is 2.42×10^{-3} . So, larger value of $J_{CP} \approx 2.65 \times 10^{-3}$ is possible and it can be probed through upcoming base-line experiments. From the Eq. (27) we can find the expression for $\sin \delta_D$. Using expressions for θ_{13} from Eq. (26), θ_{23} from Eq. (25), J_{CP} from Eq. (29) and $\cos^2 \phi$ from Eq. (22) into Eq. (27) we can have κ dependent function for $\sin \delta_D$. We have plotted δ_D with respect to κ in Fig.6 for the best fit values of θ_{12} , Δm_{32}^2 and Δm_{21}^2 . The figure reflects the prediction of δ_D as $\delta_D = 3.6^\circ$ for $\theta_{23} = 48^\circ$ at $\kappa = -2.39$.

Now we are going to see the behavior of mass eigenvalues and their sum with respect κ and hence the other observables. Using the the expressions for a^2 from Eq. (23), ϵ' from Eq. (21) and $\cos^2 \phi$ from Eq. (22) into Eq. (17) we get the κ dependent functions for mass eigen values. We have plotted m_1 , m_2 , m_3 and their sum in Fig.7 with respect to κ for the best




Figure 5: Plot of J_{CP} with respect to κ and θ_{12} . We keep Δm_{32}^2 and Δm_{21}^2 to their best fit values.




Figure 6: Plot of Dirac phase δ_D with respect to κ for the best fit values of θ_{12} , Δm^2_{32} and Δm^2_{21} .

fit values of θ_{12} , Δm_{32}^2 and Δm_{21}^2 . The observations of the plots in Fig.7 suggest that mass pattern is normal-hierarchical. We also have seen that $0.07 \text{ eV} < m_1 + m_2 + m_3 < 0.076 \text{ eV}$ in the course of variation of κ $-2.39 < \kappa < -2$ ($48^\circ > \theta_{23} > 45.8^\circ$). It also satisfy the cosmological bound $m_1 + m_2 + m_3 < 0.7 \text{ eV}$ [6].

Again parameter responsible for the $\beta\beta_{0\nu}$ experiment is also studied in the present model and the relevant quantity:

$$\begin{aligned} |(M_\nu)_{ee}| &= \left| a + \frac{2d \exp^{i\phi}}{3} \right| \\ &= a \left[1 + \frac{4\kappa(\kappa + 3) \cos^2 \phi}{9} \right]^{1/2}. \end{aligned} \quad (30)$$

Using the expressions for a and $\cos^2 \phi$ from Eq. (23) and Eq. (22) respectively, we get $|(M_\nu)_{ee}|$ in terms of model parameter κ and other physical observables. Keeping experimental values of the θ_{12} , Δm_{32}^2 and Δm_{21}^2 to their best fit value, we have plotted $|(M_\nu)_{ee}|$ with respect to κ . For the physical region of κ Fig.7 shows that $|(M_\nu)_{ee}|$ is well below the experimental bound 0.89 eV .

A point to be noted as our analysis of mixing angles are matching with [4] in the real limit ($\phi = 0^\circ$). However, the expressions for the mixing angles for complex case in [4] are not in exact correspondence with our result in the present work for $\kappa = -2$ limit. This is because in the complex analysis in [4], we have assumed that eigenvectors of M_ν construct U . This U has ability to diagonalize M_ν to its true eigenvalues but this U may not be unitary. It is better to diagonalize $h (M_\nu M_\nu^\dagger)$ which is hermitian and its diagonalizing matrix is unitary. But it is always not easy to do the same. In this paper we have solved the 18 equation from Eq. (15) and find out U in Eq. (16) which also is unitary keeping terms upto the order ϵ : $U^\dagger U = U U^\dagger = 1 + O(\epsilon^2)$.

In summary, we explore the parameter space of a softly broken A_4 symmetric model for different mixing angles and a model parameter κ . We expressed the two mixing angles θ_{12} and θ_{23} in terms of a single parameter κ , and constrained the parameter space for the best fit values of Δm_{32}^2 and Δm_{21}^2 . With the allowed parameter value we predict θ_{13} , $\theta_{13} \simeq 11^\circ$ (for 1σ deviation of θ_{23} and 2° deviation of θ_{12} about their best fit value). Utilising the above result, we expressed the CP violation parameter J_{CP} in terms of κ and $\sin \theta_{12}$ and explore the extent of J_{CP} allowed in the present model. A comparatively larger value of J_{CP} is allowed by the present model ($J_{\text{CP}} = 2.65 \times 10^{-3}$ for 1σ variation of the angle θ_{23}) and consistent with other neutrino experimental results.




Figure 7: Plot of neutrino masses m_1 , m_2 , m_3 and their sum, and also $|(M_\nu)_{ee}|$ with respect to κ for the best fit values of θ_{12} , Δm_{32}^2 and Δm_{21}^2 .

References

- [1] E. Ma and G. Rajasekaran, Phys. Rev. **D 64**, 113012 (2001); E. Ma, Mod. Phys. Lett. **A 17**, 627 (2002), K. S. Babu, E. Ma, and J. W. F. Valle, Phys. Lett. **B 552**, 207 (2003), E. Ma, Phys. Rev. **D 70**, 031901 (2004), E. Ma, Phys. Rev. **D 72**, 037301 (2005). G. Altarelli and F. Feruglio, Nucl. Phys. **B 720**, 64 (2005); See also K. S. Babu and X.-G. He, hep-ph/0507217.
- [2] G. Altarelli and F. Feruglio, Nucl. Phys. **B 741**, 215 (2006) [arXiv:hep-ph/0512103].
- [3] P. F. Harrison, D. H. Perkins, and W. G. Scott, Phys. Lett. **B 530**, 167 (2002); P. F. Harrison and W. G. Scott, arXiv: hep-ph/0402006.
- [4] B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma, M.K. Parida, Phys.Lett. **B 638** 345, 2006, hep-ph/0603059.
- [5] G. C. Branco, T. Morozumi, B. M. Nobre, M. N. Rebelo, hep-ph/0107164, hep-ph/0202036, R. N. Mohapatra and W. Rodejohann, hep-ph/0507312.
- [6] D. N. Spergel *et al.* [WMAP Collaboration], “First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Astrophys. J. Suppl. **148**, 175 (2003) [arXiv:astro-ph/0302209].